

Peptone B

RM51

Intended use

A general purpose peptone.

Contents

See pack label.

Storage and shelf life

All dehydrated culture media containers should be kept tightly closed and stored in a dry place at 10 to 25°C until the expiry date shown on the pack label.

Precautions

For *in vitro* diagnostic use only. Observe approved hazard precautions and aseptic techniques. To be used only by adequately trained and qualified laboratory personnel. Sterilise all biohazard waste before disposal. Refer to Product Safety Data sheet (available on request or via MAST[®] website).

Since the late 1980s MAST[®] has sourced all its animalderived culture media ingredients from non-bovine animals wherever possible. All animal-derived materials, including bovine materials, are from BSE-free regions of the world, are from animals certified as disease-free by qualified veterinarians and have been heat-treated in accordance with European regulations. Despite these precautions, MAST[®] Media Raw Materials must not be used in the manufacture of vaccines or food ingredients, or in the manufacture of any other high-risk products involving culture processes such as those destined for *in-vivo* or agricultural use.

Materials required but not provided

Standard microbiological supplies and equipment such as loops, MAST[®] selective supplements, swabs, applicator sticks, incinerators and incubators, etc., as well as serological and biochemical reagents and additives such as blood.

Description

A highly nutritious all purpose peptone intended for use with other culture media components. While it is not possible to produce one peptone to meet all requirements of bacteriology, MAST[®] Peptone B, has been developed to meet, in one peptone, most of the requirements of the clinical bacteriology laboratory.

It is a balanced peptone, the performance of which is constant from batch to batch. It gives good growth of fastidious organisms such as streptococci, pneumococci and Haemophilus spp., and is free from fermentable carbohydrates.

The specification of Peptone B, is similar to Peptone A, except that it has not been neutralised. It may, therefore be necessary when using this peptone to filter the prepared medium to obtain a clear product.

Mast Diagnostica GmbH Feldstrasse 20 DE-23858 Reinfeld Germany Te1: + 49 (0) 4533 2007 0 Fax: + 49 (0) 4533 2007 68 email: mast@mast-diagnostica.de Web: www.mast-group.com

Mast Diagnostic

12 rue Jean-Jacques Mention CS91106, 80011 Amiens, CEDEX 1 France Té1: + 33 (0) 3 22 80 80 67 Fax: + 33 (0) 3 22 80 99 22 email: info@mast-diagnostic.fr Web: www.mast-group.com



Procedure

Peptone B should be added at the appropriate concentration to culture media with additional ingredients e.g. agar and meat extracts as specified in the formulation being prepared.

Quality control

Check for signs of deterioration. Quality control must be performed with at least one organism to demonstrate expected performance. Do not use the product if the result with the control organism is incorrect. The list below illustrates a range of performance control strains which the end user can easily obtain.

Test Organisms	Result
Escherichia coli ATCC® 25922	Growth*
Staphylococcus aureus ATCC® 25923	Growth*

*1% w/v sterile solution of MAST[®] Peptone B with 0.5% w/v sodium chloride.

References

Bibliography available on request.