

## Brain Heart Infusion Agar

**DM104**

### Intended Use

A versatile medium for the culture of fastidious organisms.

### Contents

See pack label.

### Formulation\*

Material:	Concentration in medium:
Special peptone mixture	12.0g/litre
Yeast extract	2.0g/litre
Brain heart infusion solids	3.5g/litre
Pancreatic digest of casein	10.0g/litre
Dextrose	2.0g/litre
Sodium chloride	5.0g/litre
Disodium phosphate	0.6g/litre
Agar	12.0g/litre
Final pH: 7.3 ± 0.2	

### Storage and shelf life

All dehydrated culture media containers should be kept tightly closed and stored in a dry place at 10 to 25°C until the expiry date shown on the pack label.

### Precautions

For *in vitro* diagnostic use only. Observe approved hazard precautions and aseptic techniques. To be used only by adequately trained and qualified laboratory personnel. Sterilise all biohazard waste before disposal. Refer to Product Safety Data sheet (available on request or via MAST® website).

### Materials required but not provided

Standard microbiological supplies and equipment such as loops, MAST® selective supplements, swabs, applicator sticks, incinerators and incubators, etc., as well as serological and biochemical reagents and additives such as blood.

### Procedure

- Suspend by swirling 47.0g of powder in 1 litre of distilled or deionised water.
- Autoclave at 121°C (15 p.s.i.) for 15 minutes.
- Cool to 50 to 55°C and hold at this temperature in a water bath. Add 5 to 7% sterile defibrinated horse or sheep blood where required. Heated blood agar (chocolate) can also be prepared. Alternative growth supplements can be used.
- If required the medium can be made selective by the addition of various MAST® selective supplements.
- Pour culture plates (15 to 20ml per plate) and allow to set.
- Prepared culture plates may be used immediately or stored in plastic bags at 2 to 8°C for up to one week before use.

- Inoculate plates by surface plating, streaking out for single colonies.
- Incubate plates aerobically for 18 to 24 hours at 35 to 37°C (or alternative temperatures according to the methodology followed).

### Interpretation of results

After incubation record growth of organisms. Typical characteristics to note include colony size and morphology, pigmentation and haemolysis on blood containing medium.

### Quality control

Check for signs of deterioration. Quality control must be performed with at least one organism to demonstrate expected performance. Do not use the product if the result with the control organism is incorrect. The list below illustrates a range of performance control strains which the end user can easily obtain.

Test Organisms	Result
<i>Escherichia coli</i> ATCC® 25922	Growth, grey colonies
<i>Enterococcus faecalis</i> ATCC® 29212	Growth, grey/green colonies
<i>Pseudomonas aeruginosa</i> ATCC® 27853	Growth, grey colonies
<i>Staphylococcus aureus</i> ATCC® 25923	Growth, white/golden colonies

### References

Bibliography available on request.